


Testing methods for airway hyperresponsiveness in asthma¹⁻³



AMP, adenosine monophosphate; EVH, eucapnic voluntary hyperventilation

1. Chapman DG, Irvin CG. Clin Exp Allergy 2015;45:706–719; 2. Comberiati P, et al. Immunol Allergy Clin North Am 2018;38:545–571;

3. Cockcroft DW. Chest 2010;138(Suppl. 2):18S–24S; 4. Coates AL, et al. Eur Respir J 2017;49:1601526

Airway hyperresponsiveness and airway inflammation

Airway hyperresponsiveness to an indirect bronchial challenge test, such as mannitol, mimics active airway inflammation^{1,2}




Image copyright: Celeste Porsbjerg

This is an illustrative representation of the airway and the effect of an indirect airway hyperresponsiveness challenge; the airway and inflammatory cells are not to scale

1. Brannan JD, Loughheed MD. *Front Physiol* 2012;3:460; 2. Sverrild A, et al. *Clin Exp Allergy* 2016;46:288–297

Methacholine and mannitol challenge tests

Test	Mechanism	Measurements	Cutoff level	Interpretation of airway hyperresponsiveness
Methacholine challenge^{1*†}	Inhaled methacholine mimics the neurotransmitter acetylcholine to directly interact with muscarinic receptors on airway smooth muscle, resulting in bronchoconstriction	PC ₂₀ or PD ₂₀ to methacholine: the provoking concentration or delivered dose of methacholine required to induce 20% reduction in FEV ₁ from baseline [†]	25–100 µg / 0.13–0.50 µmol / 1–4 mg/mL	Mild
			6–25 µg / 0.03–0.13 µmol / 0.25–1 mg/mL	Moderate
			<6 µg / <0.03 µmol / <0.25 mg/mL	Marked
Mannitol challenge^{2–5}	Inhalation of mannitol rapidly increases the osmolarity of the airway surface liquid, causing stimulation of inflammatory cells (ie mast cells and eosinophils) and release of mediators, mimicking airway inflammation	PD ₁₅ to mannitol: the provoking cumulative total dose of mannitol required to induce ≥15% reduction in FEV ₁ from baseline or a 10% decrease in FEV ₁ between two consecutive mannitol doses	>155 mg	Mild indirect
			>35 mg to ≤155 mg	Moderate indirect
			≤35 mg	Severe indirect


*Challenge test results expressed as provocative dose (PD) or provocative concentration (PC) are dependent on the output rate of the administration device, time of aerosol inhalation and particle size distribution. Evidence shows that the methacholine dose, expressed as PD₂₀, allows more consistent correlation of results than PC₂₀ when comparing responses captured by different protocols. Consistency of timing between steps and from dosing to spirometry remains important to properly compare results. The PD₂₀ is the dose of methacholine that causes a 20% fall in FEV₁ and is calculated in the same way as the PC₂₀.[†] To determine changes in airway reactivity following therapy in patients known to have asthma, using doubling doses will give more precise PD₂₀ values to compare¹

FEV₁, forced expiratory volume in 1 second

1. Coates AL, et al. Eur Respir J 2017;49:1601526; 2. Hallstrand TS, et al. Eur Respir J 2018;52:1801033; 3. Comberiati P, et al. Immunol Allergy Clin North Am 2018;38:545–571;
4. Brannan JD, Loughheed MD. Front Physiol 2012;3:460; 5. Sverrild A, et al. Clin Exp Allergy 2016;46:288–297

Interpreting results of airway hyperresponsiveness bronchoprovocation tests

Change in FEV₁ as measured by PC₂₀ value by methacholine challenge test in patients with or without asthma^{1,2}



- Airway hyperresponsiveness is a valuable tool in the clinical assessment of patients with possible asthma, asthma-like symptoms, or generally normal/non-diagnostic lung function³
- In patients with asthma, bronchoconstriction (PC₂₀ as seen by reduction in FEV₁)* starts at a lower inhaled concentration of the agonist methacholine² **A**
- Maximal bronchoconstrictor responses are also greater in those with asthma^{1,2†} **B**
- Patients with **normal healthy** airways achieve a plateau response to the bronchoconstrictor stimulus, whereas patients with **mild to severe** asthma may not^{1,2}

Figure adapted from O'Byrne PM, Inman MD. Chest 2003;123(Suppl. 3):411S–416S and Nair P. J Allergy Clin Immunol Pract 2017;5:649–659

*PC₂₀ greater than 16 mg/mL represents normal airway responsiveness;⁴ 4–16 mg/mL is borderline airway hyperresponsiveness;⁴ 1–4 mg/mL represents mild airway hyperresponsiveness;⁴ 0.25–1 mg/mL represents moderate airway hyperresponsiveness;⁴ and <0.25 mg/mL represents marked airway hyperresponsiveness.⁴ †For safety reasons, methacholine testing is stopped if there has been a >20% fall in FEV₁⁴

FEV₁, forced expiratory volume in 1 second; PC₂₀, provocation concentration of methacholine causing a 20% fall in FEV₁

1. O'Byrne PM, Inman MD. Chest 2003;123(Suppl. 3):411S–416S; 2. Nair P. J Allergy Clin Immunol Pract 2017;5:649–659; 3. Cockcroft DW, et al. Allergy Asthma Clin Immunol 2020;16:14;

4. Coates AL, et al. Eur Respir J 2017;49:1601526

Contraindications for airway hyperresponsiveness testing

There are several **general contraindications** for performing tests for airway hyperresponsiveness¹

- Moderate/severe airflow limitation
- Cardiovascular problems
- Eye surgery
- Pregnancy or nursing mothers
- Inability to reproduce quality spirometry results consistently

There are also several **specific contraindications** for performing direct and indirect tests^{1,2}

Direct tests¹

- Current use of cholinesterase inhibitor medication

Indirect tests²

- Hypersensitivity to mannitol
- Presence of comorbidities that could be exacerbated by frequent coughing